



To: Prof. Noraldin Al-Deri
Early Intervention Center for Autism and
Developmental Disorders
510, 224 wasfi al tal
11953 Amman
Jordan

Homeopathy Plan

Date: 23-07-2024

Step: 1st step

| | Item name | Ingredient | Purpose | Quantity | Duration | Details |
|---|-------------------------------------------------------------------------------------|----------------------|-----------------------------|-------------------------------------------------------------------------------------|-------------|-----------------------------------|
| 1 | Swanson, Alpha Lipoic Acid, 50 mg | Alpha Lipoic Acid | White matter Development | Capsule every other day mornings. Dissolve in water, juice, honey | 4 months | Can be purchased from iherb |
| 2 | 21st Century, Standardized Ginseng Extract, 60 Vegetarian Capsule | Multiple | Behaviour improvement | Capsule every day | 4 months | Can be purchased from iherb |

ADDITIONAL INFORMATION

- A revaluation is needed after three months to reassess the development on the Ai
- Start the second step of the protocol after three months
- This assessment was developed, and its performance was validated, by the Ai model of NEURAZON and according to the findings in the report attached. The US Food and Drug Administration (FDA) has determined that clearance or approval of this method is not necessary and thus neither have been obtained. This test has been developed for clinical purposes. All test results are reviewed, interpreted and reported by our scientific and medical experts. Please refer to attached report for details.
- The provided recommendations include safe, non-drug, and natural treatments that are not a substitute for rehabilitation and training sessions. These recommendations should be followed under the supervision of a specialist. Improvement depends on the child, the underlying causes, and their response, with recommendations adjusted based on periodic tests and evaluations every three months. If any allergic symptoms appear, treatment should be stopped immediately, and the doctor should be informed to adjust the treatment plan.

